

The biology and non-chemical control of Field Penny-cress (*Thlaspi arvense* L.)

W. Bond & R.J. Turner

HDRA, Ryton Organic Gardens, Coventry, CV8 3LG, UK

Field Penny-cress

(Boor's mustard, bowyer's mustard, churl's mustard, dish mustard, mithridate mustard, treaclewort)

***Thlaspi arvense* L.**

Occurrence

Annual or overwintering plant, doubtfully native, sometimes plentiful on arable land also frequent on disturbed ground, roadsides, waste ground (Clapham et al, 1987). It is scattered over most of the UK but commoner in England (Stace, 1997). Frequent in S & E England scarcer in W & N (Rich, 1991). In a preliminary survey of arable weeds in the 1970s it was absent or rare in most of the areas studied but was common to frequent in 11% of them (Chancellor, 1977). It is not recorded above 1000 ft in the UK (Salisbury, 1961). Early and late flowering forms have been reported (Best & McIntyre, 1975). It is able to adapt to a range of environmental conditions and succeeds in both dry and wet habitats.

The seeds contain a glucocide that breaks down to form a mustard oil that can cause poisoning in stock (Best & McIntyre, 1975). Germination was inhibited in lettuce and cress seed treated with an extract of pulverised *T. arvensis* seeds (Wagenvoort & Van Opstal, 1979). The plant smells strongly when crushed and taints the milk of cows that eat it..

Biology

An annual or overwintering weed forming a compact vegetative rosette, field penny cress flowers from (March) May to October and is self pollinated. Seedlings that emerge in the autumn overwinter as a rosette and flower early in the growing season. Stem elongation occurs with the onset of flowering (Best & McIntyre, 1975). The fruit is compressed and broadly winged. There are 5-8 seeds in each of the 2 locules (Best & McIntyre, 1975), 3-8 seeds per locule (Rich, 1991). It has approximately 12 seeds per pod and 20,000 seeds per plant (Long, 1938). Salisbury (1961) gives the average seeds per pod as 16 and the average seeds per plant as 2000. Stevens (1932) gives the average seeds per plant as 7,040, Stevens (1957) quotes 900 to 2000 seeds per plant while Best & McIntyre (1975) suggest a single plant will produce 1,600 to 15,000 seeds. The range of 1000 seed weights given by different authors varied from 0.785 to 1.750 g (Stevens, 1932). Plant size and branching habit increase and flowering is delayed by increasing N (Best & McIntyre, 1975). Seed production is also likely to increase. Flowering is hastened by increasing temperatures. Spring seedlings may flower within 30-50 days of emergence and produce seed by early July.

Seed is shed over several weeks (Best & McIntyre, 1975). It is said that fresh seed will germinate in the light with adequate moisture and alternating temperatures. In lab tests, there was good germination at 10-25°C alternating temperatures in the light. Scarification of the seed coat promotes germination. Seeds can become viable 6 days after flowering (Baskin & Baskin, 1989). In the USA, plants can behave as either summer or winter annuals. Plants from seeds that germinate in autumn overwinter as

rosettes and flower in spring. Plants from seeds that germinate in spring, flower and set seed later the same year. Seeds can after-ripen at low temperature and seeds produced in autumn after-ripen over the winter and germinate in early spring. Soil disturbance increases the level of germination because of light stimulation. In the USA, freshly matured seeds in May are dormant for a short period until after-ripening has occurred in summer. Seeds that are non-dormant in autumn become conditionally dormant or dormant at low winter temperatures.

In the laboratory, stored seed did not germinate under constant temperature with or without light. Even with alternating temperatures there was less than 10% germination (Wagenvoort & Van Opstal, 1979). Germination reached 50% when the seeds were treated with fertilizer solution and alternating temperatures, and following stratification for 2 days at 5°C the level of germination increased further. Seed dormancy was overcome by chilling at between 5 & 15°C but light was still needed (Hartmann *et al.*, 1996). Without light, seeds gradually returned to a dormant state. The presence of nitrate was also important in making light stimulation more effective. It is suggested that spring cultivations should be made in unfertilised soil and that fertiliser application should be left until after crop and weed emergence.

Seed sown in a 7.5 cm layer of soil in cylinders sunk in the field and stirred periodically emerged from February to October with the main flush from February to June (Roberts, 1964). Seeds mature and shed before grain harvest. Substantial numbers germinated in the first 4 years from seed sown in trays of soil in the field but most emerged in year 1 (Chepil, 1946a). When seeds in trays of soil in the field were given different cultural treatments the highest percentage of germination was from seeds left on the soil surface (Chepil, 1946b). The deeper seeds were buried, the lower the emergence of seedlings and the greater the number of seeds that survived to the end of the experiment. Periodic cultivations increased seedling emergence. In a sandy loam soil, field seedlings emerged from the top 50 mm of soil with most coming from the top 30 mm (Unpublished information).

Persistence and Spread

Seed has a long dormancy period in excess of 3 years (Chepil, 1946a). Seeds remain viable in soil for at least 8 years (Salisbury, 1961). Seeds have survived up to 10 years in soil and still given 87% germination. In cultivated soil though, few seeds survive longer than 6 years (Best & McIntyre, 1975). In Duvel's burial experiment, seed buried at 8, 22 and 42 inches gave 11, 8 and 12% germination respectively after 1 year, 34, 52 and 8% after 6 years, 0, 11 and 1% after 30 years and none after 39 years (Toole, 1946).

Seeds will float in water for 24 hours (Best & McIntyre, 1975). Dispersal is also by wind. Viable seeds have been found in pigeon droppings and seedlings have been raised from various bird excreta (Salisbury, 1961). Apparently viable seeds were found in samples of cow manure (Mt Pleasant & Schlather, 1994). Ensilage for 8 weeks, rumen digestion in cattle for 24 hrs or a combination of both seemed to kill field penny-cress seed but some still appeared viable (Blackshaw & Rode, 1991). With rumen digestion, there was a gradual loss of viability to 30% after 24 hrs in one test but in another 98% of seeds were still able to germinate after rumen digestion. After 2 weeks of windrow composting at temperatures of 50-65 °C, penny-cress seeds were all were killed (Tompkins *et al.*, 1998).

Management

Field penny-cress seedlings should be hoed off in root crops (Long, 1938). Germination should be encouraged by surface cultivation. Do not plough in seed bearing plants. A 3-4 year ley will choke out the weed.

Acknowledgement

This review was compiled as part of the Organic Weed Management Project, OF 0315, funded by DEFRA.

References

- Baskin J M & Baskin C C** (1989). Role of temperature in regulating timing of germination in soil seed reserves of *Thlaspi arvense* L. *Weed Research* **29**, 317-326.
- Best K F & McIntyre G I** (1975). The Biology of Canadian Weeds 9. *Thlaspi arvense* L. *Canadian Journal of Plant Science* **55**, 279-292.
- Blackshaw R E & Rode L M** (1991). Effect of ensiling and rumen digestion by cattle on weed seed viability. *Weed Science* **39**, 104-108.
- Chancellor R J** (1977). A preliminary survey of arable weeds in Britain. *Weed Research* **17**, 283-287.
- Chepil W S** (1946a). Germination of weed seeds I. Longevity, periodicity of germination and vitality of seeds in cultivated soil. *Scientific Agriculture* **26** (7), 307-346.
- Chepil W S** (1946b). Germination of weed seeds II. The influence of tillage treatment on germination. *Scientific Agriculture* **26** (8), 347-357.
- Clapham A R, Tutin T G, Moore D M** (1987). *Flora of the British Isles*, 3rd edition, Cambridge University Press, Cambridge, UK.
- Hartmann K M, Mollwo A, Trautnitz P** (1996). Strategies of dormancy and germination in field penny-cress, *Thlaspi arvense* L. *Z. PflKrankh PflSchutz*, **15**, 113-123.
- Long H C** (1938). Weeds of arable land. *MAFF Bulletin* **108**, 2nd edition. HMSO, London, UK.
- Mt Pleasant J & Schlather K J** (1994). Incidence of weed seed in cow (*Bos* sp.) manure and its importance as a weed source for cropland. *Weed Technology* **8**, 304-310.
- Rich T C G** (1991). Crucifers of Great Britain and Ireland. *BSBI Handbook No. 6*. Botanical Society of the British Isles, London, UK.
- Roberts H A** (1964). Emergence and longevity in cultivated soil of seeds of some annual weeds. *Weed Research* **4** (4), 296-307.
- Salisbury E J** (1961). *Weeds & Aliens*. New Naturalist Series, Collins, London.
- Stace C** (1997). *New Flora of the British Isles*. 2nd edition. Cambridge University Press, Cambridge, UK.
- Stevens O A** (1932). The number and weight of seeds produced by weeds. *American Journal of Botany* **19**, 784-794.
- Stevens O A** (1957). Weight of seeds and numbers per plant. *Weeds* **5**, 46-55.
- Tomkins D K, Chaw D, Abiola A T** (1998). Effect of windrow composting on weed seed germination and viability. *Compost Science & Utilization* **6** (1), 30-34.

- Toole E H** (1946). Final results of the Duvel buried seed experiment. *Journal of Agricultural Research* **72** (6), 201-210.
- Wagenvoort W A & Van Opstal N A** (1979). The effect of constant and alternating temperatures, rinsing, stratification and fertilizer on germination of some weed species. *Scientia Horticulturae* **10**, 15-20.